



SAFEGUARD[®] 2.27 G/LB. CRUMBLES

PRODUCT CODE: 9100SGD

FORM: Crumble

MEDICATION:
Fenbendazole 4540 g/ton

PACKAGE SIZE: 25 lb. bag

GUARANTEED ANALYSIS:

Crude Protein (Min.)..... 15.00%
Crude Fat (Min.) 2.00%
Crude Fiber (Max.) 12.00%

INGREDIENTS:

Processed Grain By-Products, Plant Protein Products, Forage Products.

BEEF & DAIRY:

APPLICATION: For the removal and control of lungworms (*Dictyocaulus viviparus*); barberpole worms (*Haemonchus contortus*); brown stomach worms (*Ostertagia ostertagi*); small stomach worms (*Trichostrongylus axei*); hookworms (*Bunostomum phlebotomum*); thread-necked intestinal worms (*Nematodirus helvetianus*); small intestinal worms (*Cooperia punctata* and *Cooperia oncophora*); bankrupt worms (*Trichostrongylus colubriformis*); and nodular worms (*Oesophagostomum radiatum*).

Beef & Dairy Limitations: Feed as sole ration or topdress for 1 day. Retreatment may be needed after 4-6 weeks. Cattle must not be slaughtered within 13 day following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Cattle Feeding Directions:

Cattle Weight, lbs.	Lb. of Dewormer per Head	# of Head Treated per Package
400	0.4	62.5
600	0.6	41.6
800	0.8	31.2
1000	1.0	25.0
1200	1.2	20.8
1400	1.4	17.8

Feed at 2.27mg/lb. of bodyweight; 1 pound per 1000 pounds of bodyweight.

SWINE:

APPLICATION: For the removal of: Lungworms: (*Metastrongylus apri*, *Metastrongylus pudendotectus*). Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); small stomach (*Hyostromylus rubidus*); Adult and larvae (L2, L3, L4 stages – intestinal mucosal forms) whipworm (*Trichuris suis*). Kidneyworm: Adult and larvae *Stephanurus dentatus*.

Swine Limitations: Feed as sole ration in complete feed for a period of from 3-12 days to provide a total intake of 9 mg/kg (4.08 mg/lb.) of bodyweight. No withdrawal time needed.

Swine Feeding Directions:

Swine Weight, lbs.	Intake lbs.	# Days Fed	Wormer (25 lb. bags/ton of complete feed)
50	4.8	3	0.5
100	4.8	3	1
150	4.3	5	1
200	3.6	8	1

Feed at 4.08 mg/lb. of bodyweight; 1 pound per 550 pounds of bodyweight.

PRODUCT REFERENCE